

## Using a structured treatment, Friends for Life, in Norwegian outpatient clinics: results from a pilot study

Kristin D. Martinsen\*, Marianne Aalberg, Martina Gere and Simon-Peter Neumer

*Centre for Child and Adolescent Mental Health, Eastern and Southern Norway, Oslo, Norway*

*Received 27 June 2008; Accepted 17 December 2008*

**Abstract.** CBT treatment programmes for children and adolescents with anxiety disorders are promising as indicated by efficacy research. Replicating this research in ordinary clinical care is crucial in order to establish the validity of these results and disseminate empirically based treatments to practitioners. This paper presents the first experiences of using a structured programme, the Friends for Life manual, in ordinary clinical care in Norway targeting anxiety problems in children aged between 7 and 12 years. The effects of the treatment are presented as multiple single-case studies. Clinical meaningful change is considered from two perspectives; diagnostic change and changes in self-report measures. At a statistical significance level the treatment effect can be characterized as modest. Independent of the symptom reduction, the children, families and therapist are in the main satisfied with the structured approach indicating the acceptability of the programme.

**Key words:** Anxiety, clinically meaningful change, multiple single case, ordinary clinical care, treatment.

### Introduction

Anxiety disorders in youth have received considerable research attention over the past two decades. This research has confirmed the high prevalence rates (Kashani & Orvaschel, 1990), the often chronic course of the disorders (Last *et al.* 1987), that high comorbidity is the rule rather than the exception (Wittchen *et al.* 1998) and also that anxiety disorders may often be a precursor to other difficulties such as depression and substance abuse (Kendall *et al.* 2004).

The literature indicates that anxiety may increase the risk of the child not participating in age-related activities, may result in reduced academic functioning, low self-esteem, poor peer relations and family problems (Ollendick & King, 1994; Costello & Angold, 1995). Increased focus on this group of children and early intervention is thus essential given the high prevalence and bleak prognoses if left untreated (Last *et al.* 1996).

Over the last 20 years treatment programmes based on cognitive behaviour therapy for children and adolescents with anxiety disorders have been developed (Barrett, 2004;

---

\* Author for correspondence: Mrs K. D. Martinsen, Centre for Child and Adolescent Mental Health, Eastern and Southern Norway, Gullhaug Torg 4b, Oslo 0484, Norway. (email: kristin.martinsen@r-bup.no)

Kendall & Hedtke, 2006) and extensively evaluated (Barrett *et al.* 1996, 2001; Kendall & Southam-Gerow, 1996; Kendall *et al.* 2008). These treatments are today characterized as probably effective (Hibbs, 2001). While such programmes are extensively used in many countries, structured treatment programmes for emotional disorders have not been available in Norway until recently. However, the research related to the programmes is often based on efficacy studies where the conditions are close to ideal, using highly motivated therapists and children that might not be recruited from the ordinary referral system. A general concern is therefore that ordinary clinicians in outpatient clinics meeting children with multiple problems will not obtain the same positive results (Weisz *et al.* 2006). To bridge the gap between efficacy research and ordinary clinical care it is thus important to study the usefulness of such interventions in local settings with attention given to the effects across diverse patients, therapists and settings as suggested by the APA Task Force (2006). To increase the use of evidence-based treatments in ordinary clinical care, research also has to be informed by and sensitive to issues raised by practitioners (Kendall & Beidas, 2007).

The Friends programme has been translated and adapted to the Norwegian culture. In this pilot study the programme has been introduced in three outpatient clinics in Norway and this paper will focus on the effect of treatment and alternative ways of measuring this effect in addition to traditional statistical analysis. As statistically significant change may not give information about how the treatment has impacted on the client's life in terms of better quality of life, this paper will also focus on clinical meaningful change (Jacobsen & Truax, 1991). Two approaches will be used for this purpose; changes in severity rating (CRS) at diagnostic level and the Reliable Change Index (RCI; Jacobsen & Truax, 1991) on changes in self-report measures.

Last, we discuss the acceptability of the structured treatment among the families and therapists in this pilot study.

## **Method**

### ***Participants and procedure***

Children ( $n = 18$ ) aged 7–12 years (mean = 9.81 years, 48.4% girls) participated in the study. All were ethnic Norwegian. The children were recruited through the ordinary referral system of three different outpatient clinics in Southern and Eastern Norway. All children ( $n = 111$ ) in the targeted age range (7–12 years) referred to the clinics in the spring of 2006 were screened using the Children's Behavior Checklist (CBCL; Achenbach, 1991). Thirty-two children showing elevated levels of internalizing problems (T score >65) were assessed with child and parent measures.

Inclusion criteria were principal diagnosis of separation anxiety disorder (SAD), generalized anxiety disorder (GAD) or social phobia (SP) according to DSM-IV criteria (APA, 1994). The children and the parents were assessed separately to assess for anxiety disorders. Diagnosis was based on the results from the Anxiety Disorders Interview Schedule conducted with the parents (ADIS-P; Silverman & Albano, 1996). Diagnosis at post-treatment was established by a blind rater.

Exclusion criteria were IQ <70. Twenty-one of the 32 children meeting the criteria of elevated scores of internalizing problems also met the inclusion criteria of GAD, SP or SAD. Eleven children were thus excluded. Families who declined to participate in the study were

offered treatment as usual. As these 21 children were treated at three clinics, 1–2 intervention group(s) were formed at each clinic and no control condition could be established. The groups varied in size with a range from 4 to 7 children. Three children were lost to post-measurement due to high anxiety levels in the family, family problems and having started medication for attention deficit hyperactivity disorder during treatment. Final analysis was thus performed on 18 children.

The children completed a mean number of 9.2 sessions, with two booster sessions provided 3 and 4 months after the programme ended. Two children had only one booster session and two children did not attend booster sessions. There were three dropouts; after sessions 1, 2 and 5, respectively.

### **Measures**

A number of measures were used in the study. The measures included in this paper focus on the symptoms and diagnoses of the children pre- and post-intervention.

#### *Diagnosis*

The ADIS-C/P is a structured interview to diagnose anxiety disorders in children and also screen for other common DSM-IV diagnoses. Severity (CSR) is rated on a Likert scale from 0 to 8 where a rating of 8 is considered most serious and 4 is the cut-off for diagnosing the disorder. The reliability of the symptom scales of SAD, SP, specific phobia and GAD has been characterized as 'excellent' (Silverman *et al.* 2001).

#### *Child self-reports*

The Multidimensional Anxiety Scale for Children (MASC; March, 1997) is a 39-item measure of anxiety in children aged 8–19 years. The questions cover four areas; physical symptoms, avoidance, social anxiety and separation anxiety/panic. Test–retest reliability is satisfactory and found to be between 0.7 and 0.9 on the different subscales (March *et al.* 1997). The MASC has been shown to differentiate children with anxiety from children with other disorders (March, 1997). The Recent Mood and Feelings Questionnaire (MFQ; Angold, 1989) measures symptoms of depression in children and adolescents. Thirty-four items are measured on a 3-point scale. The MFQ has acceptable reliability and is a satisfactory screen for depressive disorders in children (Wood *et al.* 1995).

#### *Parent reports*

Parents reported their child's internalizing and externalizing problems on the CBCL (Achenbach, 1991). The CBCL has been extensively evaluated and has excellent psychometric properties.

The parents completed a Norwegian version of a user-satisfaction questionnaire, Fragebögen zur Beurteilung der Behandlung (FBB; Matthejat & Remschmidt, 1998) after the treatment. The scale goes from 0 to 1.5 (not satisfied), 1.5–2.5 (partly satisfied), 2.5–3.5 (largely satisfied) to 3.5–4.0 (completely satisfied). FBB has good psychometric properties.

**Table 1.** *Anxiety and severity present at T1 (n = 18)*

Diagnosis	No. of children	Severity (mean)
Generalized anxiety disorder	8	5.6
Overanxious disorder	1	6.0
Separation anxiety disorder	11	6.5
Social phobia	10	6.0
Specific phobia	6	6.2
Post-traumatic stress disorder	2	6.5

### ***Treatment materials***

A Norwegian version of the Friends for Life programme (Barrett, 2004) was used as a treatment protocol in the pilot study. The treatment is conducted in a group format and the parents are included at the end of every session. The programme is based on three major components; Learning/behaviour, Cognitive, and Physiological, and skills are taught in all areas. These skills are summarized in the name of the programme, FRIENDS, which serves as an acronym, helping the children remember the skills they learn. The programme is structured with 10 group sessions and two booster sessions. Homework activities are given after each session, and the children use workbooks throughout the programme while the group leaders are provided with a treatment manual.

Two psychologists in the project were trained and licensed as Friends facilitators at Paula Barrett's clinic: Pathways Research Centre. Psychologists from the project team ran the group sessions together with clinicians from the outpatient clinic acting as co-therapists. These clinicians participated in a 2-day training in the programme followed by monthly supervision meetings during the study.

### **Results**

#### ***Diagnostic status, severity, comorbidity and change in diagnostic status***

Table 1 indicates that many children (83%) met the criteria for more than one anxiety disorder at T1, with a mean severity rating varying from 5.6 to 6.5. SAD was the most frequent diagnosis found in the sample.

Taking all disorders into consideration, the most common comorbid disorder other than anxiety was depression/dysthemia with a mean severity rating of 5.4. Of the 18 children included, three scored over the clinical range ( $T > 64$ ) on the externalizing scale of the CBCL.

Investigations of changes in diagnostic status from pre- to post-treatment yielded a mixed picture. We found ~33% reduction in number of diagnoses at T2. SAD and depression showed the greatest absolute change with fewer children meeting the diagnostic criteria at T2. Four children no longer met the criteria for SAD and depression after treatment (see Table 2).

#### ***Clinical meaningful change at diagnosis level-severity rating (CRS)***

The mean severity rating for the group ( $n = 18$ ) at T1 was 6.15. This was only slightly reduced at T2 with a mean of 5.84.

**Table 2.** Change in diagnostic status from pre- to post-treatment (n = 18)

Diagnosis	T1 (n)	T2 (n)
Generalized anxiety disorder	8	6
Overanxious disorder	1	1
Separation anxiety disorder	11	7
Social phobia	10	8
Specific phobia	6	5
Post-traumatic stress disorder	2	1
Depression/dysthymia	5	1
Sum	43	29

**Table 3.** Severity rating pre- and post treatment for children still meeting diagnostic criteria at T2

Diagnosis	Children (n = 15)	Severity rating	
		T1	T2
Generalized anxiety disorder	6	5.50	5.33
Overanxious disorder	1	6.00	6.00
Separation anxiety disorder	7	6.86	5.57
Social phobia	8	6.57	5.29
Specific phobia	5	6.25	6.25
Post-traumatic stress disorder	1	7.00	7.00
Depression	1	8.00	8.00

At T2 the number of children meeting the criteria for any disorder was reduced from 18 to 15 as three children no longer fulfilled the criteria for any diagnosis. However, while 15 children still met the criteria for the diagnosis, the severity rating may have been reduced at T2, which may constitute clinically meaningful change. However, for most of the disorders there was no or only a modest reduction in severity rating at T2 (see Table 3). The exception was for SAD where pairwise *t* tests indicated a significant reduction in mean severity rating [ $t(6) = 2.714, p < 0.05$ ] from pre- (mean = 6.86, S.D. = 0.90) to post-treatment (mean = 5.57, S.D. = 1.51). SP showed a rather strong reduction in severity rating from 6.57 to 5.29, but with no significance [ $t(6) = 1.996, p = 0.93$ ].

For depression as a comorbid condition the severity rating at T1 ranged from 5 to 8 with a mean of 5.5. As four of the five children no longer met the diagnostic criteria for depression at T2, severity rating for only one child remained at T2, being unchanged at 8.

### ***Clinical meaningful change using self-report – RCI***

Earlier, clinical significant change has been studied with regard to the change in severity status of the children at diagnosis level. The change may also be reflected in self-report measures. *t* tests showed no significant changes in the group from pre- to post-treatment concerning anxiety symptoms [MASC:  $t(17) = 0.400, p = 0.694$ ] or depression symptoms [MFQ:  $t(16) = -0.731, p = 4.75$ ]. The RCI is a statistical approach for measuring individual change in

**Table 4.** RCI for self-report measures MASC and MFQ

Persons ( <i>n</i> = 18)	RCI – MASC total score	RCI – MFQ total score
1	–2.25	–0.23
2	–2.1	0.34
3	1.5	–0.34
4	0.3	–1.15
5	0.6	0.46
6	3	–0.46
7	–3.9	0
8	–0.45	1.83
9	–3.15	–0.57
10	–3.9	
11	0.3	–0.23
12	–0.45	0.57
13	–0.75	0.8
14	Incomplete self	Report data
15	1.35	0
16	1.5	0
17	1.5	0.34
18	–2.1	0.92

RCI, Reliable Change Index; MASC, Multidimensional Anxiety Scale for Children; MFQ, Mood and Feelings Questionnaire.

RCI  $\geq 1.96$  indicates significant change; negative sign = symptom reduction = positive effect. Reliability values used for the calculations of MASC total score: test–retest reliability,  $r = 0.933$  (March, 1997), MFQ total score: test–retest reliability,  $r = 0.80$  (Sund *et al.* 2001).

psychological disorders assessed by self-report. If the change found by RCI exceeds what can be expected based on measurement error alone, we have a statistical approach to measure clinically significant change.

All the children ( $n = 18$ ) reported their symptoms of anxiety and depression pre- and post-treatment. RCI was calculated for 17 children, as one child was excluded due to incomplete data, regardless of diagnostic status.

The results in Table 4 indicate that six of the 18 children reported a significant reduction in their anxiety symptoms and one child (number 6) showed a negative change. None of the changes in self-reported depression symptoms were significant.

### ***User satisfaction – FBB***

Another important feature of the study was to evaluate user satisfaction with treatment. The FBB measurement was used for this purpose and was scored by the parents. Most of the parents rated the programme positively with 12 of the parents rating the treatment as ‘largely satisfied’ or ‘completely satisfied’ with a mean score of 3.

However, user satisfaction was rated on two dimensions; Successfulness of treatment and Treatment process. Treatment process indicates if the parents considered that their children were appropriately treated. All parents rated Treatment process as ‘largely satisfied’ or ‘completely satisfied’ with a mean score of 3.4. In general, Successfulness of treatment was rated lower with only four parents reporting ‘largely satisfied’ or ‘completely satisfied’ and 10 rating ‘not satisfied’ or ‘partly satisfied’ on this measure. The mean score here was 2.2.

### ***Evaluation of a structured intervention by the clinicians***

All clinicians at the outpatient clinics participating in the study were asked to evaluate the usefulness of the intervention in ordinary clinical care and were invited to discuss their experiences at a 1-day conference. All the therapists wanted to continue using the structured treatment. They reported that using a high-quality structured manual was useful when treating children with anxiety disorders. Conducting treatment in a group format was also evaluated positively by the clinicians and as a possible economical way of treating these children. Moreover, it was put forward that the clinics had gained more knowledge regarding both methodology and screening measures.

### **Discussion**

The aims of the pilot study were to evaluate a structured treatment manual in ordinary clinical care and to test the acceptability of the programme.

The children included in the study had high levels of comorbidity and most met the diagnostic criteria for more than one anxiety disorder, and also had other disorders such as depression and externalizing problems. This range of difficulties was included as comorbid conditions are typical in ordinary clinical settings. At a diagnostic level the largest change observed was for SAD where 7/11 children no longer qualified for the diagnosis after treatment. For depression, 4/5 children no longer met the diagnostic criteria at T2.

Clinical meaningful change was considered by looking at changes in severity rating for the children still having the disorder at T2 and by using a statistical approach, the RCI, for evaluating clinically meaningful change at a self-reported symptom level. For the disorders SAD and SP in particular we found a substantial reduction in severity rating at T2 and for SAD the change was also statistically significant. Statistical analysis of self-report measures further indicated that 1/3 of the children showed clinically meaningful change at T2. None of the changes in self-reported depression were significant which is surprising as change in depression was noted at the diagnostic level. However, diagnosis is based on information from the parents, and discrepancy between parent information and child self-report is often reported in the literature (Kazdin, 1986).

The effects found in this pilot study may indicate that the results found in previous studies in university clinics are difficult to replicate in ordinary clinics. However, this study does have several limitations which may explain the low effect observed.

The acceptability of the programme indicates that structured approaches are welcome in ordinary clinical practice. Most of the parents were satisfied with the treatment in general. They considered that their children were treated in an appropriate way (treatment process) but were less satisfied with the successfulness of the treatment. This finding is understandable given the relatively high number of children still having symptoms of anxiety after treatment.

**Limitations**

Undertaking treatment research in Norway involves certain methodological challenges, and the present study has several limitations. The size of the outpatient clinics and the catchment area for recruiting children in Norway are relatively small and it was challenging to recruit enough children to establish a control condition. The children in the study had a wide range of comorbid problems, including externalizing problems, a condition not targeted by the programme. This may have contributed to the lack of response to the programme.

**Implications for future studies**

Despite the limitations and challenges in conducting outcome research in ordinary clinical care in Norway, this pilot study is the first to evaluate the Friends for Life programme in Norway and has provided important knowledge for future research. Given the small treatment effects, different actions could be taken to improve the effectiveness of the programme in a future study. One possibility might be to increase the specificity in the groups. This could be done by limiting the range of comorbidity and possibly having fewer anxiety disorders represented. The intensity in the programme might be increased by including more exposure in the sessions as is done in other programmes. The dosage could also be increased by having longer or more sessions and thus more time to cover the content of the programme. For some of the anxiety disorders, e.g. SP, an individual format could also be considered as being more beneficial than a group situation.

Furthermore, in order to assess ‘meaningful change’ future studies could benefit from including measures that better capture this idea. One possibility could be to operationalize each child’s treatment goal in behavioural terms and have the parents assess these pre- and post-treatment.

Given the difficulty of recruiting the necessary number of children into the study, future research on structured interventions in a group format will probably have to include multiple sites. These sites would ideally be located in close proximity so that treatment groups could be conducted at one site with participants recruited from a larger catchment area.

As most therapists in the present study were positive towards the treatment approach, it is likely that ordinary clinics could find it beneficial to participate in a larger study. Training of the therapists both with regard to the programme, and assessment as well as providing continuous supervision is essential to ensure the quality of the diagnostic process and fidelity to the programme used. However, this might be a challenge with regard to the ecological validity of the results.

Addressing these limitations and dilemmas in future studies may enable us to evaluate structured interventions in ordinary clinical care more fully and thus contribute to the effectiveness research which eventually may be crucial in bridging the gap between research and practice.

**Summary of main points**

- Structured treatments are welcome in ordinary clinical care.
- Effect of treatment may be measured using alternative approaches.

- Clinical meaningful change may be a useful measure of bridging the gap between research and clinic.
- Conducting treatment research in ordinary clinics presents challenges with regard to recruiting a sufficient number of children and thus establishing control conditions.

### Declaration of Interest

None.

### Recommended follow-up reading

- Kendall PC, Beidas RS** (2007). Smoothing the trail for dissemination of evidence based practice for youth: flexibility within fidelity. *Professional Psychology: Research and Practice* **38**, 13–20.
- Weisz JR, Jensen-Doss A, Hawley K** (2006). Evidence-based youth psychotherapies versus usual clinical care. A meta-analysis of direct comparisons. *American Psychologist* **61**, 671–689.

### References

- Achenbach T** (1991). *Manual for the Child Behavior Checklist and 1991 Profile*. University of Vermont, Department of Psychiatry.
- APA** (1994). *Diagnostic and Statistical Manual of Mental Disorders*, 4th edn (DSM-IV). Washington, DC: American Psychiatric Association.
- Angold A** (1989). Structured assessments of psychopathology in children and adolescents. In: *The Instruments of Psychiatric Research* (ed. C. Thompson), pp. 271–303. Chichester: John Wiley & Sons.
- APA Task Force** (2006). APA Presidential Task Force on Evidence Based Practice. *American Psychologist* **61**, 271–285.
- Barrett PM** (2004). *Friends for Life: Group Leader's Manual*. Brisbane, Australia: Australian Academic Press.
- Barrett PM, Dadds MR, Rapee RM** (2001). Cognitive-behavioral treatment of anxiety disorders in children: long-term (6-year) follow up. *Journal of Counseling and Clinical Psychology* **69**, 135–141.
- Barrett PM, Dadds MR, Rapee RM** (1996). Family treatment of childhood anxiety: a controlled trial. *Journal of Counseling and Clinical Psychology* **64**, 333–342.
- Costello EJ, Angold A** (1995). Epidemiology. In: *Anxiety Disorders in Children and Adolescent* (ed. J. S. March), pp. 109–124. New York: Guilford Press.
- Hibbs ED** (2001). Evaluating empirically based psychotherapy research for children and adolescents. *European Child and Adolescent Psychiatry* **10**, 1/3–1/11.
- Jacobsen NS, Truax P** (1991). Clinical significance: a statistical approach to defining Meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology* **59**, 12–19.
- Kashani JH, Orvaschel H** (1990). A community study of anxiety in children and adolescents. *American Journal of Psychiatry* **147**, 313–318.
- Kazdin AE** (1986). Developing effective treatments for children and adolescents. In: *Psychosocial Treatments for Child and Adolescent Disorders – Empirically Based Strategies for Clinical Practice* (ed. E. D. Hibbs and P. S. Jensen), pp. 9–18. Washington, DC: American Psychological Association.
- Kendall PC, Beidas RS** (2007). Smoothing the trail for dissemination of evidence based practice for youth: flexibility within fidelity. *Professional Psychology: Research and Practice* **38**, 13–20.
- Kendall PC, Hudson JL, Gosch E, Flannery-Schroeder E** (2008). Cognitive-behavioral therapy for anxiety disordered youth: a randomized clinical trial evaluating child and family modalities. *Journal of Consulting and Clinical Psychology* **76**, 282–297.

- Kendall PC, Hedtke KA** (2006). *Cognitive-Behavioral Therapy for Anxious Children: Therapist Manual*. Ardmore, PA: Workbook Publishing.
- Kendall PC, Safford S, Flannery-Schroeder E, Webb A** (2004). Child anxiety treatment: Maintenance of outcomes in adolescence and impact on substance use and depression at 7.4 year follow-up. *Journal of Consulting and Clinical Psychology* **72**, 276–287.
- Kendall PC, Southam-Gerow MA** (1996). Long term follow up of a cognitive-behavioral therapy for anxiety-disorders in youth. *Journal of Consulting & Clinical Psychology* **64**, 724–730.
- Last CG, Hersen M, Kazdin AE, Finkelstein R, Strauss CC** (1987). Comparison of DSM-III separation anxiety and overanxious disorders: Demographic characteristics and patterns of comorbidity. *Journal of American Academy of Child and Adolescent Psychiatry* **26**, 527–531.
- Last CG, Perrin S, Hersen M, Kazdin AE** (1996). A prospective study of childhood anxiety disorders. *Journal of the American Academy of Child and Adolescent Psychiatry* **35**, 1502–1510.
- March JS** (1997). *MASC – Multidimensional Anxiety Scale for Children – Technical Manual*. Toronto: Multi-Health Systems Inc.
- March JS, Parker J, Sullivan K, Stallings P, Connors CK** (1997). The Multidimensional Anxiety Scale for Children (MASC): factor structure, reliability, and validity. *Journal of the American Academy of Child and Adolescent Psychiatry* **36**, 554–565.
- Mattejat F, Remschmidt H** (1998). *Fragebogen zur Berteilung der Behandlung (FBB)*. Göttingen: Hogrefe.
- Ollendick TH, King NJ** (1994). Empirically supported treatments for children and adolescents: current status. In: *Child and Adolescent Therapy: Cognitive-Behavioral Procedures*, 3rd edn (ed. P. C. Kendall), pp. 492–520. New York: Guilford Press.
- Silverman WK, Albano AM** (1996). *Anxiety Disorders Interview Schedule for DSM-IV – Parent version*. San Antonio, TX: Graywind Publications.
- Silverman WK, Saavedra LM, Pina AA** (2001). Test–retest reliability of anxiety symptoms and diagnosis with the Anxiety Disorders Interview Schedule for DSM-IV: Child and Parent versions. *Journal of the American Academy of Child and Adolescents Psychiatry* **40**, 937–944.
- Sund AM, Larsson B, Wichstrøm L** (2001). Depressive symptoms among young Norwegian adolescents as measured by The Mood and Feelings Questionnaire (MFQ). *European Child & Adolescent Psychiatry* **10**, 222–229.
- Weisz JR, Jensen-Doss A, Hawley KM** (2006). Evidence-based youth psychotherapies versus usual clinical care. A meta-analysis of direct comparisons. *American Psychologist* **61**, 671–689.
- Wittchen H-U, Nelson CB, Lachner G** (1998). Prevalence of mental disorders and psychosocial impairments in adolescents and young adults. *Psychological Medicine* **28**, 109–126.
- Wood A, Kroll L, Moore A, Harrington R** (1995). Properties of the Mood and Feelings Questionnaire in adolescent psychiatric outpatients: a research note. *Journal of Child Psychology and Psychiatry* **36**, 327–334.

### Learning objectives

- Considering alternative ways of measuring effect of treatment – clinical meaningful change.
- Evaluating usefulness of a structured treatment programme for clients, parents and practitioners.